

## Key Points

- The Centers for Disease Control and Prevention (CDC) has published provisional life expectancy estimates stating that the average life expectancy of Americans decreased by a year and a half during 2020 primarily due to COVID-19. This statement can be misleading due to the life expectancy measure used by the CDC.
- The CDC uses “period” life expectancy, which assumes that the COVID-19 mortality experience of 2020 would continue for every year in the future. While this measure may be useful for year-over-year comparisons in normal times, it tends to exaggerate the effect of nonrecurring events.
- “Cohort” life expectancy is how most people think about life expectancy, as it is an estimate of the actual number of years that an individual may live, reflecting expected future conditions. The temporary impacts of increased deaths due to COVID-19 in 2020 and into 2021 will have a much smaller effect on cohort life expectancy, as it is assumed that those effects will lessen over time.
- The Actuaries Longevity Illustrator can help individuals estimate how long they might live. Actuaries are well positioned to help bring clarity to understanding of longevity and life expectancy.



AMERICAN ACADEMY of ACTUARIES

*Objective. Independent. Effective.™*

1850 M Street NW, Suite 300  
Washington, DC 20036

202-223-8196 | [www.actuary.org](http://www.actuary.org)

Craig Hanna, Director of Public Policy  
Linda K. Stone, Senior Pension Fellow

© 2021 American Academy of Actuaries. All rights reserved.

# Interpreting Pandemic-Related Decreases in Life Expectancy

DECEMBER 2021

The Centers for Disease Control and Prevention (“CDC”) has published provisional life expectancy estimates in its Vital Statistics Surveillance Report<sup>1</sup> that the average life expectancy of Americans decreased by a year and a half during 2020, primarily due to the impact of COVID-19.<sup>2</sup> This development understandably raises concerns because it creates an impression that the average American will live a year and a half less than they otherwise would have due to COVID-19.

But what is “life expectancy”—and does the CDC’s statement really mean what many people think it means?

The CDC report, which is based on deaths during 2020, garnered widespread attention that Americans had lost a year and a half of life expectancy due to the pandemic. A reduction of roughly three years in life expectancy was cited for Black and Hispanic Americans. This decrease in life expectancy follows several years of smaller life expectancy declines (on average), which have run counter to the long-term trend of increases in longevity.

While these changes in life expectancy might seem dramatic, it can be difficult to interpret what this data means. The notion of life expectancy can be confusing because there are different measures of life expectancy that are used in different contexts. It is important to fully understand these measures to interpret the CDC analysis of the data for 2020. The CDC focuses on a hypothetical measure known as “period” life expectancy. While the CDC report does distinguish this measure from the “cohort” life expectancy measure that aligns with the more intuitive understanding of the metric, described below, the CDC’s focus is on changes in period life expectancy that

<sup>1</sup> *Report No. 015*; Centers for Disease Control and Prevention; July 2021.

<sup>2</sup> CDC publication *Mortality in the United States, 2020* includes final statistics for 2020 showing a 1.8-year decline in life expectancy at birth.

were largely driven by the increase in deaths attributable to COVID-19. As discussed below, this focus on period life expectancy can create confusion.

**Period life expectancy**, measured as of a given age, is the remaining number of years that a hypothetical individual can expect to live (on average) if rates of death prevailing during a given period continue indefinitely. When mortality rates increase significantly during a given year—as has been the case during the COVID-19 pandemic—period life expectancy will decline noticeably because it is calculated as if these higher rates of mortality will continue in all future years. This often-cited measure of life expectancy can vary dramatically from one year to the next. The CDC report highlights period life expectancy at birth—the number of years that a newborn would be expected to live if the current pandemic conditions persist throughout the newborn’s entire lifetime.

**Cohort life expectancy** (also known as “generation” life expectancy) considers changes in rates of mortality that are expected to occur in the future. A pandemic or other significant event that has a substantial effect on short-term population mortality but is expected to subside will have a much smaller effect on cohort life expectancy than it does on period life expectancy.

Period life expectancy may be a useful metric for comparing general population mortality experience for different periods of time. Nonetheless, cohort life expectancy is likely what most people envision when thinking about the concept of life expectancy because cohort life expectancy is an estimate of the actual number of years that a typical individual might be expected to live based on reasonable expectations for future conditions. For this reason, cohort life expectancy is the measure used by the Actuaries Longevity Illustrator.<sup>3</sup> Both measures of life expectancy are based on probabilities and capture the average expectation of a large population. A given individual may live many years beyond that average or may die well short of this average expected life span.

This issue brief discusses these concepts, how life expectancies are calculated, how to interpret them, and the limitations of these measures. In developing these concepts, the underlying concepts of mortality rates, mortality (or “life”) tables, and the data used to calculate mortality rates and life expectancy are discussed.

<sup>3</sup> [Actuaries Longevity Illustrator](#); American Academy of Actuaries and Society of Actuaries.

The Pension Committee, which authored this issue brief, includes Elena Black, MAAA, FSA, FCA, EA—*Chairperson*; Grace Lattyak, MAAA, FSA, FCA, EA—*Vice Chairperson*; Michael Antoine, MAAA, FSA, FCA, EA; Michael Bain, MAAA, ASA, FCA, FSPA, EA; Rachel Barnes, MAAA, FSA, FCA, CERA, EA; Margaret Berger, MAAA, FSA, FCA, EA; Maria Carnovale, MAAA, FSA, EA; Jonathan de Lutio, MAAA, EA; C. David Gustafson, MAAA, FCA, EA; Scott Hittner, MAAA, FSA, FCA, EA; Lloyd Katz, MAAA, FSA, FCA, EA; Maria Kirilenko, MAAA, ASA, FCA, EA; Tonya Manning, MAAA, FSA, FCA, EA; Maria Moliterno, MAAA, ASA, EA; A. Donald Morgan, MAAA, FSA, FCA, EA; James Ritchie, MAAA, ASA, FCA, EA; Mark Shemtob, MAAA, FSA, FCA, EA; Mary Stone, MAAA, FSA, FCA, EA; Todd Tauzer, MAAA, FSA, FCA, CERA; Hal Tepfer, MAAA, FSA, FCA, MSPA, EA;

The committee gratefully acknowledges the contributions of Immediate Past Chairperson Bruce Cadenhead, MAAA, FSA, FCA, EA

## Mortality rates and mortality tables

In the course of their work, actuaries consider and study uncertain events and make assumptions about probabilities of these events occurring. For example, the probability of a person of a given age dying within the next year is what is known as an “age-specific mortality rate.” Another way to look at this is that the estimated likelihood of a person living to the next age is 100% minus this age-specific mortality rate.

In general, a mortality table is composed of age-specific mortality rates for a range of ages, usually from birth to the oldest age to which anyone is expected to survive. These tables are also sometimes called “life tables.”<sup>4</sup> While many actuaries use mortality tables for different populations in the course of their work, life expectancies are not directly used in actuarial practice; rather, they are a separate output of the underlying mortality tables.

## Factors that affect mortality rates

Mortality tables are created by tabulating observed mortality experience data from large populations. It is well established that different populations may experience very different survivorship expectations, based on the population’s characteristics. Many factors can affect mortality rates, several of which are highlighted in this section.

First, by their very construction, mortality tables recognize the obvious factor of age.

Another well-established and well-known factor is gender because, as a group, females tend to live longer than males. Most mortality tables are constructed showing distinct probabilities of death for males and for females.<sup>5</sup>

Other factors that may be reflected by constructing separate tables or by developing other adjustments<sup>6</sup> include socioeconomic status, level of education, race, occupation, health status including body mass index, level of physical activity, smoking status, other lifestyle-related factors, and geography.

<sup>4</sup> For example, the CDC [publishes](#) mortality tables for U.S. population, under the name “Life Tables.”

<sup>5</sup> Although an increasing number of people identify as nonbinary, there is not yet sufficient data available to develop separate rates of mortality for these individuals.

<sup>6</sup> Such adjustments may be achieved by multiplying rates by a factor, or “stepping back or forward” by simply realigning the existing rates with different ages.

## How mortality experience in the U.S. has changed over time

U.S. population mortality rates have generally decreased over time. As mortality rates decrease, longevity and life expectancy increase. Over the past decade, this trend of longevity increases has stalled, with life expectancy at birth virtually unchanged between 2010 and 2018,<sup>7</sup> due largely to the effects of factors such as an increased use of opioids and the prevalence of obesity offsetting the generally positive trend. Meanwhile, more affluent segments of the population have continued to experience increases in life expectancy during this period.<sup>8</sup>

Historically, gains in longevity can be tracked to simple improvements in living conditions, such as the use of clean water and waste removal. More recent factors that resulted in increases in longevity in the U.S. include advances in medicine (for example, the development of antibiotics), expanded access to health care through introduction of Medicare and Medicaid, continual and general improvements in education, health awareness, reduction in smoking rates, and improved standards of living.

Opinions vary among demographers, actuaries, academics, and other professionals who study and analyze mortality and associated trends as to whether the historic longevity improvement trend will resume and, if so, at what pace. In addition, the COVID-19 pandemic has produced at least a temporary dip in the trend of increasing longevity. The long-term consequences of this pandemic on mortality and health status are largely unknown at this time. Some speculate that long-term effects of COVID-19 could continue to increase rates of mortality relative to pre-pandemic levels, while others believe that medical advances driven by the pandemic, such as development of vaccines, could lead to greater improvements in longevity.

## The relationship between mortality rates and life expectancy

The concept of probability is commonly used to quantify unknown and uncertain events. For example, using a mortality table can help a 65-year-old evaluate their chance of celebrating their next birthday. Computing a life expectancy is a statistical process of “collapsing” a series of age-specific survival probabilities into one number that represents the expected number of additional years a person might live.<sup>9</sup>

---

<sup>7</sup> “Changes in Life Expectancy at Birth, 2010–2018”; Centers for Disease Control and Prevention; Jan. 30, 2020

<sup>8</sup> *U.S. Population Mortality Observations—Updated with 2019 Experience*; Society of Actuaries; January 2021.

<sup>9</sup> In other words, this single measure is a sum of the probabilities, derived from the mortality table, of a person of a given age to live to all future ages.

The mathematics behind the calculation of life expectancy uses the concept of “expected value” of a person’s future lifetime. The word “expected” communicates that the computed result is not a *fact*; it’s what is *expected on average*. For example, if an actuary tells you that a 40-year-old has a life expectancy of 41.7 years,<sup>10</sup> it does not mean that the 40-year-old will live to age 81.7 exactly. Some individuals, age 40 today, will die before that age while others will live past that age. But if we tracked the future lifetimes of 100,000 people who are 40 years old today, we would *expect* that upon the recorded death of the last of the group, these 100,000 people will have lived *an average* of 41.7 additional years.

Another issue that may cause confusion is that life expectancy measures for U.S. populations are often reported for the population as a whole, blending many characteristics that influence mortality into a single reported measure. Life expectancy can be reported for a combined population of males and females, covering individuals living in different parts of the country, and reflecting different socioeconomic characteristics.

Yet another complicating feature of this measure is that when reporting life expectancies, many—including the CDC in the report cited above—illustrate these values as of different starting ages. It is common to track life expectancy at birth, as well as at other ages. And, because (for example) a 40-year-old person has already survived from birth to age 40, the 40-year-old’s expected age at death will be higher than the expected age at death for a newborn (or for anyone who is younger than 40), because the life expectancy for a anyone younger than age 40 must also incorporate the possibility that they may not survive to age 40.<sup>11</sup>

The recent attention to the significant decreases in life expectancy reported by CDC focused on the “at birth” period life expectancy measure, but this nuanced distinction may be lost on the general public. When a 65-year-old person hears a “1.5-year decrease in life expectancy,” they may not realize that the reported drop from 78.8 years to 77.3 refers to the “at birth” measure. For a 65-year-old, the CDC’s recent report shows a drop in the period life expectancy of 0.8 years—from 19.6 in 2019<sup>12</sup> to 18.8 in 2020.<sup>13</sup>

10 Using Social Security [Life Expectancy Calculator](#) for a male, age 40 today, utilizing “cohort” life expectancy methodology and Social Security Administration (SSA) assumptions for future male mortality.

11 The term “expected age at death” is used here to avoid confusion over the term “life expectancy,” as it applies to individuals who have already lived a number of years. For example, for a 40-year-old person, the CDC report shows a period life expectancy of an additional 39.6 years, resulting in an expected age at death of 79.6. For newborns, on the other hand, the life expectancy and the expected age at death are the same (77.3).

12 See CDC publication [Mortality in the United States, 2019](#).

13 Note, for example, as reported by CDC for 2019, at birth the life expectancy was 78.8 while for a 65-year-old it was 19.6, resulting in average expected age at death of 84.6—which exceeds 78.8 by 5.8 years—because the 65-year-old already survived the first 65 years of their life.

## Cohort and period life expectancies

The group of 40-year-old people discussed in the previous example is a “cohort” because they were all born in the same year. Computing the life expectancy for a cohort can be done by tracking the group from the time analysis is begun (in this case, age 40) and continuing as the number of years each member of this cohort lives is recorded until the last death of the group is recorded.

If actuaries had a way to peer into the future so that they could observe the actual ages at death for each cohort member, the cohort life expectancy for this group could be calculated as the average of observed life spans. For historical cohorts, such as for those born in 1880 who were alive and age 40 in 1920, average life span may be calculated from historical records.

For a cohort of current 40-year-old people, actuaries routinely use actuarially developed mortality tables to determine the number of years members of this group are expected to live. As discussed earlier, the mortality table is typically developed from experience for a relevant population for a particular period. In using this table to determine life expectancy, this approach assumes that the mortality rates measured in this period apply to all future years.

To estimate period life expectancy in the recent report, the CDC used mortality rates for the general U.S. population for 2020. This means someone who is 40 years old in 2020 is assumed to experience these mortality rates in all future years. In 25 years, for example, when this person is 65, they are assumed to have the same mortality experience as someone who is age 65 in 2020. This methodology of life expectancy calculation is a standard approach that the CDC uses when reporting life expectancy figures, and which is widely cited.

To develop an estimate more aligned with a commonly accepted understanding of how long people alive today are expected to live, this standard approach is refined by incorporating mortality improvement or changes over time into this calculation. Any such projection depends upon the assumption that future mortality rates will change over time. This assumption will have a significant effect on life expectancy.

Period life expectancy measures demonstrate fluctuations that reflect events that influenced mortality in this particular period.<sup>14</sup> For example, the Spanish flu pandemic of 1918 resulted in a dramatic decrease in period life expectancy, which was more than offset by an increase in period life expectancy the next year. A male baby born in 1917 had a period life expectancy of 52.2 years, while a male baby born in 1918 had a period life expectancy of only 45.3 years—a reduction of almost 7 years.<sup>15</sup> The following year, a male newborn had a period life expectancy of 54.2, an increase of almost 9 years over the period life expectancy calculated in 1918 for a newborn male. These changes are much larger than those seen thus far with COVID-19, demonstrating the relative severity of that earlier pandemic relative to the current one.

It is instructive to review the impact of calculating life expectancies on a cohort basis, rather than a period basis, for these three cohorts of male newborns in the late 1910s. Using mortality rates published by the SSA for years after 1917, for a cohort of 1917 male newborns, the average life span was 59.4; for the 1918 cohort, average life span was 60.0; and for 1919, it was 61.5. Even these differences are heavily influenced by the fact that the 1917 and 1918 cohorts had to survive the high rates of death during 1918, while the 1919 cohort did not.

If both period and cohort life expectancy are measured as of 1920 for each of these groups (the 3-year-old children who were born in 1917, 2-year-old children who were born in 1918 and 1-year-old children who were born in 1919), differences are observed in these measures as they narrow substantially because the high rates of mortality during 1918 have no effect on those who survived to 1920. This is summarized in the table below.



Year of Birth	Life Expectancy at birth		Life expectancy in 1920	
	Period	Cohort	Period	Cohort
1917	52.2	59.4	58.3	65.3
1918	45.3	60.0	58.7	66.0
1919	54.2	61.5	58.6	66.2

This table, as well as the graphs that follow, use historical and projected U.S. population mortality rates published by Social Security Administration in conjunction with the 2020 Trustees Report.

<sup>14</sup> The federal government began keeping mortality statistics going back to 1900. Data is available from the CDC and SSA going back to that year. Methodologies evolved and changed over time, and those changes as well as the passage of time impacted the results. Due to these differing methodologies, numbers may be significantly different depending on the source.

<sup>15</sup> For these numbers, [historical U.S. population mortality rates](#) published by the Social Security Administration in conjunction with the publication of the 2020 Trustees Report were used.

The graphs below show how both period and cohort life expectancy have changed over time. The significant dip in period life expectancy during 1918 and subsequent recovery during 1919 can be seen in these results, while the annual changes in cohort life expectancy are less volatile. Due to historical and expected future improvements in mortality rates, cohort life expectancy is consistently higher than period life expectancy.





## How not to interpret life expectancy

Life expectancy can be a confusing concept and is often subject to misinterpretation and misunderstanding. An individual may conclude that the expected life span they hear about or read about is a prediction of the exact number of years they can expect to live, and that the CDC's recent report means that they will not survive as long as they may have expected previously. In fact, however, anyone alive in 2021 reading about the CDC report will already have survived the high rates of mortality experienced during 2020.

## How temporary phenomena, such as a pandemic, affect life expectancy measures

The long-term effect of COVID-19 on longevity is as yet unknown. If the effects are temporary and rates of mortality return to pre-pandemic levels and continue to improve in line with historical trends, period life expectancy measures reported by the CDC should see a corresponding rebound. The COVID-19-related effects published thus far by the CDC are, as noted in this issue brief, *period* life expectancies, which assume that current conditions continue indefinitely and have thus decreased during the pandemic. If pre-COVID-19 mortality rates return, the CDC's measure of life expectancy should increase in future years. Cohort life expectancy, which is more heavily influenced by long-term expectations, will likely see at most a small change in response to any temporary increase in mortality.

The considerations discussed in this issue brief highlight the need to carefully consider the meaning of life expectancy trends, COVID-19-related or otherwise. The American Academy of Actuaries Pension Committee has produced this issue brief to provide useful information to help the reader better understand the complexities involved in the measurement of life expectancy estimates.

The American Academy of Actuaries is a 19,500-member professional association whose mission is to serve the public and the U.S. actuarial profession. For more than 50 years, the Academy has assisted public policymakers on all levels by providing leadership, objective expertise, and actuarial advice on risk and financial security issues. The Academy also sets qualification, practice, and professionalism standards for actuaries in the United States.